# *Contents Volume 89, 1994*

An AMS radiocarbon method to determine the emplacement time of recent deep-sea turbidites
J. Thomson and P.P.E. Weaver . . . . . 1
Kopara in Polynesian atolls: early stages of formation of calcareous stromatolites
C. Défarge, J. Trichet, A. Maurin and M. Hucher . . . . . 9
Tsunami deposits in a lacustrine sequence of the Sanriku coast, northeast Japan
K. Minoura, S. Nakaya and M. Uchida . . . . . 25
Persistence of topographic features as a result of non-tectonic processes
J.P. le Roux . . . . . 33
Tertiary sedimentary rocks associated with the Harcuvar core complex in Arizona (U.S.A.): insights into paleogeographic evolution during displacement along a major detachment fault system
J.C. Yarnold . . . . . 43
Evolving fluvial–transitional–marine deposition through the Cambrian sequence of Jordan
B.S. Amireh, W. Schneider and A.M. Abed . . . . . 65
Peat-accumulating depositional systems of Sarawak, East Malaysia
J.R. Staub and J.S. Esterle . . . . . 91
Tunnel-valley formation in northwest Germany—geology, mechanisms of formation and subglacial bed conditions for the Bornhöved tunnel valley
J.A. Piotrowski . . . . . 107
Geometry, palaeoenvironments and relative sea-level (accommodation space) changes in the Messinian Murdjadjo carbonate platform (Oran, western Algeria): consequences
J.-J. Cornée, J.-P. Saint Martin, G. Conesa and J. Muller . . . . . 143

*Book Review*
*The Geology of Libya, Vols. IV–VII*, by M.J. Salem et al. (Editors)—W.A. Berggren . . . . . 159

The Late Devensian and Holocene evolution of Barmouth Bay, Wales
P. Larcombe and C.F. Jago . . . . . 163
Depositional mechanisms as revealed from grain-size measures of the Palaeoproterozoic Kolhan Siliciclastics, Keonjhar District, Orissa, India
S.K. Ghosh and B.K. Chatterjee . . . . . 181
Compositional trends in the Permian sandstones from the Denison Trough, Bowen Basin, Queensland reflect changing provenance and tectonics
R. Ahmad, J.C. Tipper and R.A. Eggleton. . . . . 197
Braidplain sedimentation in a Proterozoic rift basin: the São João da Chapada Formation, southeastern Brazil
M.A. Martins-Neto . . . . . 219
Storm features of siliciclastic shelf sedimentation in the mid-Cretaceous epeiric seaway of northern Australia
A.A. Krassay. . . . . 241
Provenance of the Lower Tertiary Murree redbeds (Hazara–Kashmir Syntaxis, Pakistan) and initial rising of the Himalayas
S. Critelli and E. Garzanti . . . . . 265
Marine to fluvial transition: Proterozoic Upper Rewa Sandstone, Maihar, India
P.K. Bose and P.P. Chakraborty . . . . . 285
Late Ordovician–Early Silurian glaciofluvial deposits preserved in palaeovalleys in South Jordan
J.H. Powell, B.K. Moh'd and A. Masri . . . . . 303

A neutron goniometer study of the preferred orientation of calcite in fine-grained deep-sea carbonate
L. Ratschbacher, A. Wetzel and H.-G. Brokmeier . . . . . 315
Factors controlling the growth of stromatoporoid biostromes in the Ludlow of Gotland, Sweden
S. Kershaw and M. Keeling . . . . . 325